



King's Research Portal

DOI:

[10.1016/j.ijcard.2017.11.097](https://doi.org/10.1016/j.ijcard.2017.11.097)

Document Version

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Steinmetz, M., Broder, M., Hösch, O., Lamata, P., Kutty, S., Kowallick, J. T., Staab, W., Ritter, C. O., Hasenfuß, G., Paul, T., Lotz, J., & Schuster, A. (2018). Atrio-ventricular deformation and heart failure in Ebstein's Anomaly — A cardiovascular magnetic resonance study. *International Journal of Cardiology*.
<https://doi.org/10.1016/j.ijcard.2017.11.097>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal


Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Contents lists available at ScienceDirect

International Journal of Cardiology

journal homepage: www.elsevier.com/locate/ijcard

Atrio-ventricular deformation and heart failure in Ebstein's Anomaly – A cardiovascular magnetic resonance study

Michael Steinmetz^{a,d,*}, Marike Broder^{c,d,1}, Olga Hösch^{a,d,2}, Pablo Lamata^{f,g,1}, Shelby Kutty^{e,3}, Johannes T. Kowallick^{b,d,3}, Wieland Staab^{b,d,2}, Christian Oliver Ritter^{b,d,2}, Gerd Hasenfuß^{c,d,2}, Thomas Paul^{a,d,3}, Joachim Lotz^{b,d,2}, Andreas Schuster^{c,d,h,**,1}

^a Department of Pediatric Cardiology and Intensive Care Medicine, University Medical Center, Georg-August-University Göttingen, Heart Center, Germany

^b Inst. for Diag. and Interventional Radiology, University Medical Center, Georg-August-University Göttingen, Heart Center, Germany

^c Department of Cardiology and Pneumology, University Medical Center, Georg-August-University Göttingen, Germany

^d DZHK, German Center for Heart Research, partner site Göttingen, Germany

^e University of Nebraska Medical Center/Children's Hospital and Medical Center, Omaha, NE, USA

^f Department of Computer Science, University of Oxford, Oxford, United Kingdom

^g Division of Imaging Sciences and Biomedical Engineering, The Rayne Institute, St. Thomas' Hospital, King's College London, London, United Kingdom

^h Department of Cardiology, Royal North Shore Hospital, The Kolling Institute, Northern Clinical School, University of Sydney, Sydney, Australia

ARTICLE INFO

Article history:

Received 2 February 2017

Received in revised form 4 November 2017

Accepted 27 November 2017

Available online xxxx

Keywords:

Ebstein's Anomaly

CMR

Feature tracking

Strain

R/L-Volume-Index

Heart failure

ABSTRACT

Purpose: We aimed to quantify atrial and ventricular myocardial deformation in Ebstein's Anomaly (EA) in a case-control study with cardiovascular magnetic resonance (CMR) feature tracking and to correlate changes in cardiac performance with the severity of disease and clinical heart failure parameters.

Materials and methods: Atrial and ventricular deformation was measured using CMR feature tracking in 30 EA and 20 healthy control subjects. Atrial performance was characterized using longitudinal strain and strain rate parameters for reservoir function, conduit function and booster pump function. Ventricular performance was characterized using RV and LV global longitudinal strain (ϵ_l) and LV circumferential and radial strain (ϵ_c and ϵ_r). Volumetric measurements for the ventricles including the Total Right/Left-Volume-Index (R/L-Volume-Index) and heart failure markers (BNP, NYHA class) were also quantified.

Results: EA patients showed significantly impaired right atrial performance, which correlated with heart failure markers (NYHA, BNP, R/L-Volume-Index). LA function in EA patients was also impaired with atrial contractile function correlating with NYHA class. EA patients exhibited impaired RV myocardial deformation, also with a significant correlation with heart failure markers.

Conclusion: CMR feature tracking can be used to quantify ventricular and atrial function in a complex cardiac malformation such as EA. EA is characterized by impaired quantitative right heart atrio-ventricular deformation, which is associated with heart failure severity. While LV function remains preserved, there is also significant impairment of LA function. These quantitative performance parameters may represent early markers of cardiac deterioration of potential value in the clinical management of EA.

© 2017 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Correspondence to: M. Steinmetz, Department Pediatric Cardiology and Intensive Care Medicine, University Medical Center, Georg-August-University Göttingen, Robert-Koch-Str. 40, 37075 Göttingen, Germany.

** Correspondence to: A. Schuster, Department of Cardiology and Pneumology, University Medical Center, Georg-August-University Göttingen, Robert-Koch-Str. 40, 37099 Göttingen, Germany.

E-mail addresses: michael.steinmetz@med.uni-goettingen.de (M. Steinmetz), andreas.schuster@med.uni-goettingen.de (A. Schuster).

¹ M.S., M.B., P.L. and A.S. take responsibility for study design and planning, all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

² O.H., W.S., C.O.R., G.H., J.L. all take responsibility for scientific discussion and improvement of the study design and the manuscript.

³ S.K., J.T.K., T.P. take responsibility for study design and planning, careful revision of data interpretation, revision of the manuscript.

1. Introduction

Ebstein's Anomaly (EA) of the tricuspid valve (TV) accounts for approximately 0.5% of congenital heart disease (CHD) patients. Many EA patients suffer from progressive right heart failure and impaired physical exercise capacity at some stage in their life [1,2]. The pathophysiology comprises a dysplastic and tethered TV displaced towards the right ventricular (RV) apex with subsequent "atrialization" of the RV (aRV), higher grade tricuspid valve regurgitation (TVR) and RA and RV volume overload [3]. Additionally, deterioration of LV function is frequently observed in EA [4,5]. To date, the pathophysiological changes that ultimately cause progression to global heart failure in EA are

unknown. Moreover there is a lack of biomarkers of clear clinical value that indicate cardiac deterioration at an early stage in EA.

The clinical relevance of impaired atrial function in heart failure development is increasingly recognized [6]. Right atrial (RA) function has been studied in some congenital heart disease (CHD) [7,8] using echocardiography and CMR [9]. Both echocardiography and CMR-FT are equally able to discriminate left and right atrial phasic performance based on strain (ϵ) and strain rate (SR) imaging [10–12]. The reservoir phase, characterized by collection of pulmonary venous return during ventricular systole can be quantified by global peak longitudinal total strain (ϵ_s) and global peak positive SR (SRs). The conduit phase that describes the passage of blood to the ventricle during early ventricular diastole is indicated by global longitudinal passive strain (ϵ_e) and global peak early negative SE (SRe). The contractile phase or booster pump function describes the intensity of atrial contraction resulting in augmentation of ventricular filling during late ventricular diastole and can be quantified measuring global longitudinal active strain (ϵ_a) and global peak late-negative SR (SRa) during atrial contraction [12]. These parameters have been studied in non-symptomatic participants of the Multi-Ethnic-Study of Arteriosclerosis [13,14] demonstrating prognostic value in heart failure prediction and in patients with atrial fibrillation showing a close link to stroke occurrence.

Alterations in atrial function – which reflect changes in myocardial deformation itself but are also dependent on RV function, loading conditions and tricuspid regurgitation – may possibly represent early markers for deterioration of overall cardiac function, but have not been defined in EA [11]. Also the interplay of atrial and ventricular performance of the right and left heart may play a role in EA disease progression. Thus, we sought to assess atrial and ventricular function in EA using a comprehensive CMR protocol and myocardial FT analysis. Moreover, we aimed to correlate changes in biatrial and biventricular performance with the severity of EA and clinical parameters of heart failure.

2. Methods

2.1. Study population

From a total of fifty-eight patients with EA in the database of the Department of Pediatric and Adult CHD, University Medical Center Göttingen, Germany, 31 patients with EA were prospectively recruited in this case-control study and gave written informed consent to participate. Exclusion criteria included age < 10 years, implantable electronic cardiac devices, associated complex cardiac malformations and claustrophobia. The 31 patients (21 male) were examined prospectively within one day between January 2013 and July 2013 and compared to 20 healthy matched controls (10 male). Six of the EA patients had undergone previous cardiac surgery (2 Glenn-Anastomosis, 2 TV reconstructions, one TV replacement and one patient with both, a TV reconstruction and Glenn-Anastomosis). Seven of eight patients with atrial septum defect (ASD) had undergone ASD closure (3 interventional, 4 surgical). Sinus rhythm was documented during the examination and no higher degree AV blocks were noted. The study protocol included medical history, New York Heart Association (NYHA) classification, clinical examination, brain natriuretic peptide (BNP) and CMR. The study protocol complied with the Declaration of Helsinki and was approved by the local clinical ethics committee.

2.2. CMR imaging

CMR was performed according to current guidelines [15]. All CMR measurements were performed at a 1.5 Tesla MRT-“Symphony”-scanner (Siemens Medical Systems, Erlangen, Germany). All patients were examined according to a standardized imaging protocol for EA and without sedation. A 16 channel phased-array-receiver coil was used for signal reception. Ventricular dimensions and function were assessed using stacks of multislice-multiphase steady state free precession cine images in ventricular short-axis as well as in axial orientations. Standard imaging parameters were as follows: repetition time = 44 ms, echo time = 1.3 ms, flip angle = 20°, slice thickness = 5 mm, spatial resolution 1.6 × 1.6 mm; parallel imaging acceleration factor 2; 12 s breath hold; with retrospective gating. Pulmonary artery flow was measured using phase contrast through-plane measurements in the main pulmonary artery just above the pulmonary valve and then used to calculate the TVR with the equation introduced by Fratz et al. ($TR = \frac{(RVSV - PA_{ante})}{(RVSV - PA_{retro})} \times 100$) [16].

2.3. Feature tracking

Myocardial FT was performed using dedicated software (TomTec Imaging Systems, 2D CPA MR, Cardiac Performance Analysis, Version 1.1.2.36, Unterschleissheim, Germany). Results are based on averaging of three repeated measurements by the same observer to maximize reproducibility as previously demonstrated [17].

2.4. Atrial feature tracking analysis

The 4-chamber view showing all 4 cardiac chambers was used to calculate LA and RA strain and strain rate [18]. As the anatomy can vary significantly in EA patients this will be referred to as the horizontal long axis view in the course of this manuscript. Atrial endocardial borders were manually drawn using a point-and-click approach at the end of ventricular diastole representing the time of the lowest atrial filling. Atrial borders of the anatomical RA were tracked at the site of the true tricuspid annulus from the lateral wall to the septal RA wall. We thereby excluded the atrialized portion of the RA, which is marked by the displaced tricuspid valve and could not be tracked. After the automated tracking algorithm was applied, tracking performance was visually controlled and, in case of insufficient border tracking, manually corrected and the algorithm reapplied. In case of persisting insufficient tracking quality, the corresponding segment was excluded from the analysis.

The three main components of atrial physiology were quantified as follows (Figs. 1A and B): 1) atrial reservoir function representing the collection of venous return during ventricular systole using total strain (RA ϵ_s /LA ϵ_s) and peak positive strain rate (SRs), 2) atrial conduit function representing the passage of blood to the ventricles during early ventricular diastole using passive strain (RA ϵ_e /LA ϵ_e) and peak early negative strain rate (SRe) and 3) atrial contractile booster pump function representing the augmentation of ventricular filling during late ventricular diastole using active strain (RA ϵ_a /LA ϵ_a) and peak late negative strain rate (SRa) [6].

2.5. Ventricular analysis

Ventricular strain was calculated from horizontal long axis and short-axis slices as previously described [19]. The horizontal long axis was used to quantify endocardial RV and LV global longitudinal strain (RV ϵ_l /LV ϵ_l). In order to calculate global circumferential and radial LV short-axis strain (LV ϵ_c and LV ϵ_r), endo- and epicardial contours, were drawn on basal, mid-ventricular and apical short-axis slices. Global LV ϵ_c was calculated as average from subendocardial and subepicardial LV ϵ_c .

Volumetric analysis was performed with commercially available software (QMass, Medis, The Netherlands). For EA patients the functional right ventricle (fRV) was used for RV measurements and contours were drawn to the insertion of the displaced TV leaflets. For ventricular volumetric analysis, endocardial contours for RV and LV were drawn in all short axis slices at end systole and end diastole. Short axis orientation was employed deliberately since previous work has demonstrated similar results for short axis and transverse quantification of ventricular function [20], as well as to ensure comparability between EA patients and healthy volunteers. EF was calculated from end diastolic volume indexed to BSA (EDVi) and end systolic volume indexed to BSA (ESVi).

The Total Right/Left-Volume-Index (R/L-Volume-Index) as a clinical severity parameter of EA was calculated from end diastolic volume measurements ($R/L\text{-Volume-Index} = \frac{RA+RV+FRV}{LA+LV}$) [20].

2.6. Statistical analysis


Statistical analysis was performed using Microsoft Excel (Microsoft Corporation, Redmond, Washington, USA), Statistica (Stat Soft, North Melbourne, Australia) and IBM SPSS Statistics version 22 (IBM Corporation, Armonk, New York, USA). Data are expressed as mean (\pm standard deviation). After assessment for normality using the Kolmogorov-Smirnov test, the data were compared using a *t*-test and the Mann-Whitney-*U* test for parametric and non-parametric data, respectively. Spearman's correlation coefficients were calculated for relations between functional indexes and heart failure parameters and for relations between the functional indexes of atria and ventricles. A sub-group analysis was performed after exclusion of EA patients with previous corrective surgery (24 patients left).

Measurements were performed by two independent skilled observers. The first observer (M.B.) analysed all data based on average curves of three analyses repetitions and repeated the analyses of the functional atrial parameters in 10 randomly selected subjects after a period of 4 weeks for the intra-observer variability. The second observer (J.T.K.) performed the analysis in the same 10 subjects for the analysis of the inter-observer variability. Reproducibility was assessed using the coefficient of variation [21] and the method described by Bland and Altman [22]. Due to multiple comparisons, *p*-values for Spearman's correlation were reported for descriptive purposes. For all other comparisons, *p*-values < 0.05 were considered statistically significant.


3. Results

CMR-FT was successfully performed in 30 patients with EA and all controls. One patient with EA was excluded due to insufficient image

A



B



quality from motion artifacts. The mean age of the EA patients was 26.3 years (controls 23.7 years, $p = 0.238$).

Overall, 85.3% of the atrial segments were included in the EA group and 94.6% in the control group. Exclusion of atrial segments was predominantly associated with poor tracking quality due to insertion of pulmonary veins in the LA and poor imaging quality associated with breathing motion.

3.1. Right atrial function

RA reservoir and booster pump function were significantly reduced in the EA group as compared to controls (Table 1). Conduit function was significantly lower in EA based on SRe, whilst there was a trend towards reduction in ϵ derived conduit function (Table 1).

3.1.1. Correlation with heart failure markers

RA function exhibited correlations with heart failure parameters in EA patients (see Table 2): 1) atrial reservoir function correlated significantly with BNP level (RA ϵ s and SRs) and NYHA class ($p < 0.05$ for RA ϵ s; and a strong trend for RA SRs, $p = 0.067$); 2) atrial conduit function (RA ϵ e and SRe) correlated with the R/L-Volume Index; and 3) atrial booster pump function (RA ϵ a and SRa) correlated with NYHA class and BNP (see Table 2).

There was no significant correlation between RA deformation parameters and the degree of TVR.

3.2. Right ventricular function

RV volume was higher, and RVEF and RV longitudinal strain (RV ϵ l) were significantly lower in EA patients compared with healthy controls (see Table 1).

3.2.1. Correlation with heart failure markers

RV volume and RVEF correlated with the R/L-Volume Index. RV ϵ l correlated with all heart failure parameters (R/L-Volume Index, NYHA class, BNP) (see Table 2).

3.2.2. Atrio-ventricular coupling

Atrio-ventricular coupling analysis revealed correlations of RV ϵ l with atrial reservoir and conduit function (see Table 3).

3.3. Left atrial function

EA patients had decreased LA reservoir (ϵ s) and conduit function (ϵ e, SRe) as well as reduced booster pump function (SRa) (see Table 1).

3.3.1. Correlation with heart failure markers

LA booster pump function showed a correlation with NYHA class based on strain and SR, respectively (NYHA–LA ϵ a $r = -0.455$, $p = 0.017$ and SRa $r = 0.455$, $p = 0.017$).

3.4. Left ventricular function

There was a trend towards lower LV longitudinal strain (LV ϵ l) in EA patients compared to controls ($p = 0.067$), while LV volumetric parameters did not differ between groups.

3.4.1. Correlation with heart failure markers

LV volumetric data correlated with NYHA class and the R/L-Volume-Index (LVEF total–NYHA $r = -0.411$, $p = 0.030$ and EDVi–R/L Index $r = -0.491$, $p = 0.009$).

3.5. Impact of cardiac surgery

The correlation of heart failure stages with both reservoir function (NYHA, BNP) and booster pump function (NYHA) remained, irrespective of the exclusion of the patients who underwent corrective surgery (data not shown). Conversely, the association of RA booster pump function and BNP could not be detected when looking at the EA subgroup without previous cardiac surgery ($p > 0.05$).

Furthermore, cardiac surgery did not impact on the association of the severity of EA (R/L-Volume-Index) and RA conduit function.

3.6. Inter- and intraobserver variability

Atrial strain and SR were reproducible on an inter- and intraobserver level with low values for mean difference and coefficient of variation (range 10.5 to 32.8 depending on individual parameter, supplementary material supplied upon request).

4. Discussion

Our study describes a well-characterized, functional phenotype based on atrial and ventricular mechanical function in EA. We demonstrate that CMR-FT is a suitable technique for quantification of cardiac atrial and ventricular deformation in a complex congenital heart malformation such as EA. We can show for the first time, that in EA CMR-FT derived RA and RV function, but to some extent also LA function, was impaired and correlated with heart failure markers. Correlation analysis of RA function with heart failure markers revealed the following: 1) patients with more severe heart failure stages (expressed by BNP and NYHA class) showed larger reduction of RA reservoir function and RA booster pump function, and 2) patients with more severe EA (as expressed by the R/L-Volume-Index) had lower values for RA conduit function.

It is important to note, that these findings were independent of previous cardiac surgery and are thus applicable to a heterogeneous EA patient collective. Moreover, it is important to understand that RA function is dependent on loading conditions and RV and tricuspid valve function at the time of the exam. Changes in RA function are not necessarily reflective of overall intrinsic atrial function alone.

Besides functional impairment of the RA, LA conduit function was also affected to a lesser extent. Interestingly, whilst LA reservoir and conduit function were lower as compared to control subjects only LA booster pump function was related to heart failure severity as defined by NYHA. This is paralleled by association of reduced RA booster pump function with NYHA class in our cohort. Similar to findings in non-congenital left heart disease [23], this highlights the potential importance of the active atrial contractile function in heart failure in EA.

4.1. Right heart function

Standard RV functional parameters derived from CMR including EF and volumes do not correlate well with exercise capacity and heart failure markers in EA patients [24]. A lack of correlation between RV volumes and EF with NYHA and BNP was observed in the present study, as well, but we identified a correlation of RV strain and several RA metrics (Table 2).

Quantification of RA function in CHD is receiving increasing interest [25] and impaired RA function has been reported in congenital right heart disease Tetralogy of Fallot (TOF) [26] or ASD [27]. Reduced RA reservoir function and elevated conduit function occur in TOF with higher grade pulmonary regurgitation and large RV EDVi [26]. In our cohort of EA, RA reservoir and booster pump function were

Fig. 1. A: Right atrial feature tracking in a patient with Ebstein's Anomaly and a healthy control. B: Left atrial feature tracking in a patient with Ebstein's Anomaly and a healthy control. RA ϵ l: right atrial longitudinal strain; LA ϵ l: left atrial longitudinal strain; SR: strain rate.

Table 1
Comparison of functional parameters of EA patients and healthy controls.

	EA patients	Controls	P-value	EA patients	Controls	P-value
	Right atrium			Left atrium		
Reservoir function						
Strain (ϵ_s (%))	20.50 \pm 9.91	28.79 \pm 10.86	0.009	17.81 \pm 9.45	21.06 \pm 5.79	0.037
Strain rate (SRs (/s))	0.96 \pm 0.35	1.20 \pm 0.42	0.037	0.74 \pm 0.31	0.89 \pm 0.27	0.094
Conduit function						
Strain (ϵ_e (%))	14.31 \pm 7.93	18.29 \pm 8.85	0.094	13.14 \pm 7.95	15.28 \pm 5.30	0.040
Strain rate (SRe (/s))	−0.65 \pm 0.29	−0.90 \pm 0.42	0.021	−0.76 \pm 0.38	−1.08 \pm 0.44	0.007
Booster pump function						
Strain (ϵ_a (%))	6.19 \pm 4.88	10.49 \pm 5.27	0.006	4.67 \pm 3.38	5.78 \pm 3.76	0.300
Strain rate (SRa (/s))	−0.59 \pm 0.40	−0.90 \pm 0.48	0.020	−0.41 \pm 0.32	−0.61 \pm 0.35	0.047
	Right ventricle			Left ventricle		
EDVi (ml/m ²)	117.54 \pm 58.10	76.13 \pm 16.97	<0.001	72.35 \pm 10.42	76.60 \pm 10.19	0.172
ESVi (ml/m ²)	66.85 \pm 40.00	36.22 \pm 9.09	<0.001	29.64 \pm 6.82	28.81 \pm 8.14	0.708
EF total (%)	44.77 \pm 8.33	52.99 \pm 5.23	<0.001	59.07 \pm 7.50	62.75 \pm 7.67	0.281
Longitudinal strain (ϵ_l (%))	−13.48 \pm 6.26	−19.66 \pm 3.60	<0.001	−15.67 \pm 4.96	−18.37 \pm 4.78	0.067
Circumferential strain (ϵ_c (%))				−17.61 \pm 4.63	−17.81 \pm 3.13	0.870
Radial strain (ϵ_r (%))				25.65 \pm 9.83	28.22 \pm 9.75	0.378

P-value from t-test or Mann-Whitney-U test, bold p-values indicate statistical significance.

EA: Ebstein's Anomaly; AVI: atrial volume, indexed; EF: ejection fraction, indexed; ϵ_s : strain; SR: strain rate; EDVi: end-diastolic volume, indexed; ESVi: end-systolic volume, indexed; ϵ_c : circumferential strain; ϵ_r : radial strain.

impaired and associated with higher NYHA class and increased BNP. This is in line with observations in TOF patients [9].

The mechanism behind impaired RA deformation might be a higher degree of RA preload (determined by a) central venous return and b) TR induced volume load) and an unfavourable pre-dilation of atrial myofibres (Frank-Starling mechanism). Changes in atrial strain and strain rate, as observed in our study, are reflective of contractility and distensibility of the myofibrils, i.e. possibly alterations in the optimal Frank-Starling curve. A study of Ren et al. links changes in LA function in patients with mitral regurgitation to the Frank Starling Mechanism and atrial pre-load conditions [28]. This supports our hypothesis that loading conditions and the Frank Starling mechanism play a role in altered atrial function in EA. Since the atrialized RV contracts in synchrony with the RV and thus against a closed TV, the degree of dyssynchrony of the aRV and anatomical RA may be another contributor to changes in the atrial contraction. This remains speculative, since aRV tracking is not feasible with our technique.

RV function in EA patients was also compromised, as expressed by impaired RV ϵ_l . Changes in RV ϵ_l depict changes in RV myocardial

contraction and are – comparable to RA function – dependent on ventricular loading conditions. However, RV ϵ_l correlation with heart failure markers suggest that these changes in deformation may be clinically relevant.

Unifying the above observations, we hypothesized that cardiac dysfunction in EA is linked to decreased RA deformation – possibly based on a shift away from the optimal Frank-Starling curve – coupled with impaired RV longitudinal function (i.e. reduced RV ϵ_l in the EA group). Spearman analysis of RA–RV interaction in our cohort confirmed positive correlation of RV strain with RA reservoir and conduit function and supports this hypothesis.

4.2. Left heart function

Our results show that LA function is also affected, albeit to a lesser extent compared to the RA, while LV systolic function was normal. Even though systolic LV function may be preserved in our collective, LA reservoir and conduit function are reduced and may reflect LV diastolic dysfunction. This hypothesis is in line with Inai's description of diastolic dysfunction in EA with preserved LV EF [29].

4.3. Limitations

This was a single center cross sectional study. Our cohort only comprises a small number of patients with higher degree heart failure (NYHA III or IV), which is due to the low incidence of EA.

Table 2
Correlation of right atrial and ventricular volumes and functional indexes from MR with heart failure parameters of EA patients.

	R/L-Volume-Index	NYHA	BNP
<i>Right atrium</i>			
<i>Reservoir function</i>			
Strain (ϵ_s)	− 0.449 (0.019)	− 0.446 (0.015)	− 0.637 (<0.001)
Strain rate (SRs)	n.s.	−0.345 (0.067)	− 0.487 (0.009)
<i>Conduit function</i>			
Strain (ϵ_e)	− 0.416 (0.031)	n.s.	− 0.516 (0.005)
Strain rate (SRe)	− 0.400 (0.039)	n.s.	n.s.
<i>Booster pump function</i>			
Strain (ϵ_a)	n.s.	− 0.560 (0.002)	− 0.420 (0.026)
Strain rate (SRa)	n.s.	− 0.476 (0.009)	− 0.409 (0.031)
<i>Right ventricle</i>			
EDVi	0.844 (<0.001)	n.s.	n.s.
ESVi	0.918 (<0.001)	n.s.	n.s.
EF	− 0.670 (<0.001)	n.s.	n.s.
Strain (ϵ_l)	− 0.644 (<0.001)	− 0.466 (0.012)	− 0.395 (0.042)

Spearman's correlation coefficient (p-value), n.s.: non-significant.

RAVi: right atrial volume, indexed; EF: ejection fraction; ϵ_s : strain; SR: strain rate; EDVi: end-diastolic volume, indexed; ESVi: end-systolic volume, indexed; R/L-Volume-Index: Total Right/Left-Volume-Index; NYHA: New York Heart Association classification; BNP: brain natriuretic peptide.

Table 3
Correlation of right atrial strain parameters with right ventricular volumes and functional indexes from MR of EA patients.

	EDVi	ESVi	RV EF	RV ϵ_l
<i>Right atrium</i>				
<i>Reservoir function</i>				
Strain (ϵ_s)	n.s.	n.s.	n.s.	0.660 (<0.001)
Strain rate (SRs)	n.s.	n.s.	n.s.	0.460 (0.014)
<i>Conduit function</i>				
Strain (ϵ_e)	n.s.	n.s.	n.s.	0.518 (0.005)
Strain rate (SRe)	n.s.	n.s.	n.s.	0.474 (0.011)
<i>Booster pump function</i>				
Strain (ϵ_a)	n.s.	n.s.	n.s.	n.s.
Strain rate (SRa)	n.s.	n.s.	n.s.	n.s.

Spearman's correlation coefficient (p-value), n.s.: non-significant.

ϵ_s : strain; SR: strain rate; EDVi: end-diastolic volume, indexed; ESVi: end-systolic volume, indexed.

To discern the role of the RA conduit function further, CMR-FT analysis of both the RA and aRV would be necessary. However, since CMR-FT relies on tracking myocardial borders and the aRV lies within the fRV only separated from it by the malformed TV, there are no myocardial borders that can be tracked by CMR-FT. This makes CMR-FT analysis of the aRV technically impossible.

Finally, age-wise maturation of atrial function have been reported in the echocardiography literature [30]. Due to the rarity of the disease and the small patient numbers available, effects of age on atrial function have not been accounted for in the present study.

5. Conclusion

Right atrial and ventricular myocardial strain and strain rate are reduced in EA and can be reliably quantified using CMR-FT. CMR derived longitudinal right atrial and ventricular strain appear to be markers of heart failure in EA, since a close correlation of these parameters with NYHA class, BNP and R/L-Volume-Index was observed. Alterations in LA function were also noted. Incorporating atrial and ventricular deformation indices into CMR assessment may help to improve the management and decision making for intervention or operation in EA patients.

Abbreviations

aRV	atrialized right ventricle
ASD	atrial septum defect
BNP	brain natriuretic peptide
BSA	body surface area
CHD	congenital heart disease
CMR	cardiovascular magnetic resonance imaging
ϵ	strain
ϵ_c	circumferential strain
ϵ_l	longitudinal strain
ϵ_a	atrial active longitudinal strain
ϵ_e	atrial passive longitudinal strain
ϵ_s	atrial total longitudinal strain
ϵ_r	radial strain
EA	Ebstein's Anomaly
EDVi	end diastolic volume indexed to body surface area
EF	ejection fraction
ESVi	end systolic volume indexed to body surface area
fRV	functional right ventricle
FT	feature tracking
LA	left atrium
LV	left ventricle
NYHA	New York Heart Association
RA	right atrium
	R/L-Volume-Index Total-Right/Left-Volume-Index
RV	right ventricle
SR	strain rate
S _{Ra}	atrial peak late negative strain rate
S _{Re}	atrial peak early negative strain rate
S _{Rs}	atrial peak positive strain rate
TOF	Tetralogy of Fallot
TVR	tricuspid valve regurgitation
TV	tricuspid valve

Conflict of interest

The authors report no relationships that could be construed as a conflict of interest.

Disclosures

None.

Acknowledgements

The study was supported by the German Center for Cardiovascular Research (DZHK).

P.L. holds a Sir Henry Dale Fellowship funded jointly by the Wellcome Trust and the Royal Society under grant number 099973/Z/12/Z. This work was also supported by the Wellcome EPSRC Centre for Medical Engineering at King's College London (WT 203148/Z/16/Z).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijcard.2017.11.097>.

References

- [1] D. Tobler, S. Yalonetsky, A.M. Crean, J.T. Granton, L. Burchill, C.K. Silversides, R.M. Wald, Right heart characteristics and exercise parameters in adults with Ebstein anomaly: new perspectives from cardiac magnetic resonance imaging studies, *Int. J. Cardiol.* 165 (1) (2013) 146–150.
- [2] O. Hösch, S.C. Alt, T. Paul, J. Lotz, M. Steinmetz, A. Schuster, Managing Ebstein's anomaly of the tricuspid valve: impact of cardiovascular magnetic resonance, *J. Cardiol. Ther.* 1 (7) (2014) 154–159.
- [3] C.H. Attenhofer Jost, H.M. Connolly, J.A. Dearani, W.D. Edwards, G.K. Danielson, Ebstein's anomaly, *Circulation* 115 (2) (2006) 277–285.
- [4] P.J. Golecki, F.H. Sheehan, S.S.M. Chen, P.J. Kilner, M.A. Gatzoulis, The shape and function of the left ventricle in Ebstein's anomaly, *Int. J. Cardiol.* 171 (3) (2014) 404–412.
- [5] C.H. Attenhofer Jost, H.M. Connolly, P.W. O'Leary, C.A. Warnes, A.J. Tajik, J.B. Seward, Left heart lesions in patients with Ebstein anomaly, *Mayo Clin. Proc.* 80 (3) (2005) 361–368.
- [6] B.D. Hoit, Left atrial size and function: role in prognosis, *J. Am. Coll. Cardiol.* 63 (6) (2014) 493–505.
- [7] M. Abd El Rahman, T. Raedle-Hurst, A. Rentzsch, H.-J. Schäfers, H. Abdul-Khalik, Assessment of inter-atrial, inter-ventricular, and atrio-ventricular interactions in tetralogy of Fallot patients after surgical correction. Insights from two-dimensional speckle tracking and three-dimensional echocardiography, *Cardiol. Young* 25 (7) (2014) 1254–1262.
- [8] J. Hou, H.-K. Yu, S.J. Wong, Y.-F. Cheung, Atrial mechanics after surgical repair of tetralogy of Fallot, *Echocardiography* 32 (2014) 126–134.
- [9] S.E. Luijnenburg, R.E. Peters, R.J. van der Geest, A. Moelker, J.W. Roos-Hesselink, Y.B. de Rijke, B.J.M. Mulder, H.W. Vliegen, W.A. Helbing, Abnormal right atrial and right ventricular diastolic function relate to impaired clinical condition in patients operated for tetralogy of Fallot, *Int. J. Cardiol.* 167 (3) (2013) 833–839.
- [10] J.T. Kowallick, S. Kutty, F. Edelmann, A. Chiribiri, A. Villa, M. Steinmetz, J.M. Sohns, W. Staab, N. Bettencourt, C. Unterberg-Buchwald, G. Hasenfuß, J. Lotz, A. Schuster, Quantification of left atrial strain and strain rate using cardiovascular magnetic resonance myocardial feature tracking: a feasibility study, *J. Cardiovasc. Magn. Reson.* 16 (1) (2014) 60.
- [11] J.T. Kowallick, J. Lotz, G. Hasenfuß, A. Schuster, Left atrial physiology and pathophysiology: role of deformation imaging, *World J. Cardiol.* 7 (6) (2015) 299.
- [12] A. Schuster, K.N. Hor, J.T. Kowallick, P. Beerbaum, S. Kutty, Cardiovascular magnetic resonance myocardial feature tracking: concepts and clinical applications, *Circ. Cardiovasc. Imaging* 9 (2016) e004077.
- [13] M. Habibi, H. Chahal, A. Opdahl, O. Gjesdal, T.M. Helle-Valle, S.R. Heckbert, R. McClelland, C. Wu, S. Shea, G. Hundley, D.A. Bluemke, J.A. Lima, Association of CMR-measured LA function with heart failure development, *JACC Cardiovasc. Imaging* 7 (6) (2014) 570–579.
- [14] Y.Y. Inoue, A. Alissa, I.M. Khurram, K. Fukumoto, M. Habibi, B.A. Venkatesh, S.L. Zimmerman, S. Nazarian, R.D. Berger, H. Calkins, J.A. Lima, H. Ashikaga, Quantitative tissue-tracking cardiac magnetic resonance (CMR) of left atrial deformation and the risk of stroke in patients with atrial fibrillation, *J. Am. Heart Assoc.* 4 (4) (2015) e001844.
- [15] S. Fratz, T. Chung, G.F. Greil, M.M. Samyn, A.M. Taylor, B. Valsangiacomo, R. Emanuela, S.-J. Yoo, A.J. Powell, Guidelines and protocols for cardiovascular magnetic resonance in children and adults with congenital heart disease: SCMR expert consensus group on congenital heart disease, *J. Cardiovasc. Magn. Reson.* 15 (1) (2013) 51.
- [16] S. Fratz, C. Janello, D. Müller, M. Seligmann, C. Meierhofer, T. Schuster, C. Schreiber, S. Martinoff, J. Hess, A. Kühn, M. Vogt, H. Stern, The functional right ventricle and tricuspid regurgitation in Ebstein's anomaly, *Int. J. Cardiol.* 167 (1) (2012) 258–261.
- [17] A. Schuster, V.-C. Stahnke, C. Unterberg-Buchwald, J.T. Kowallick, P. Lamata, M. Steinmetz, S. Kutty, M. Fasshauer, W. Staab, J.M. Sohns, B. Bigalke, C. Ritter, G. Hasenfuß, P. Beerbaum, J. Lotz, Cardiovascular magnetic resonance feature-tracking assessment of myocardial mechanics: intervendor agreement and considerations regarding reproducibility, *Clin. Radiol.* 70 (2015) 989–998.
- [18] J.T. Kowallick, G. Morton, P. Lamata, R. Jogiya, S. Kutty, G. Hasenfuß, J. Lotz, E. Nagel, A. Chiribiri, A. Schuster, Quantification of atrial dynamics using cardiovascular magnetic resonance: inter-study reproducibility, *J. Cardiovasc. Magn. Reson.* 17 (2015) 36.
- [19] A. Schuster, S. Kutty, A. Padiyath, V. Parish, P. Gribben, D.A. Danford, M.R. Makowski, B. Bigalke, P. Beerbaum, E. Nagel, Cardiovascular magnetic resonance myocardial

- feature tracking detects quantitative wall motion during dobutamine stress, *J. Cardiovasc. Magn. Reson.* 13 (1) (2011) 58.
- [20] O. Hösch, J.M. Sohns, T.-T. Nguyen, P. Lauerer, C. Rosenberg, J.T. Kowallick, S. Kutty, C. Unterberg, A. Schuster, M. Faßhauer, W. Staab, T. Paul, J. Lotz, M. Steinmetz, The Total right/left-volume-index: a new and simplified cardiac magnetic resonance measure to evaluate the severity of Ebstein's anomaly of the tricuspid valve: a comparison with heart failure markers from various modalities, *Circ. Cardiovasc. Imaging* 7 (2014) 601–609.
- [21] F. Grothues, G.C. Smith, J.C.C. Moon, N.G. Bellenger, P. Collins, H.U. Klein, D.J. Pennell, Comparison of interstudy reproducibility of cardiovascular magnetic resonance with two-dimensional echocardiography in normal subjects and in patients with heart failure or left ventricular hypertrophy, *Am. J. Cardiol.* 90 (1) (2002) 29–34.
- [22] J.M. Bland, D.G. Altman, Statistical methods for assessing agreement between two methods of clinical measurement, *Lancet* 1 (8476) (1986) 307–310.
- [23] V. Melenovsky, B.A. Borlaug, B. Rosen, I. Hay, L. Ferruci, C.H. Morell, E.G. Lakatta, S.S. Najjar, D.A. Kass, Cardiovascular features of heart failure with preserved ejection fraction versus nonfailing hypertensive left ventricular hypertrophy in the urban Baltimore community: the role of atrial remodeling/dysfunction, *J. Am. Coll. Cardiol.* 49 (2) (2007) 198–207.
- [24] O. Hösch, T.-T. Nguyen, P. Lauerer, A. Schuster, S. Kutty, W. Staab, C. Unterberg-Buchwald, J.M. Sohns, T. Paul, J. Lotz, M. Steinmetz, BNP and haematological parameters are markers of severity of Ebstein's anomaly: correlation with CMR and cardiopulmonary exercise testing, *Eur. Heart J. Cardiovasc. Imaging* 16 (6) (2015) 670–675.
- [25] S. Kutty, Q. Shang, D. Danford, M. Steinmetz, A. Schuster, T. Kuehne, P.B. Beerbaum, S. Sarikouch, Abnormal right atrial performance in surgically repaired Tetralogy of Fallot: a CMR feature tracking analysis, *Int J. Cardiology* 248 (2017) 136–142.
- [26] E. Riesenkampff, N. Al-Wakeel, S. Kropf, C. Stamm, V. Alexi-Meskishvili, F. Berger, T. Kuehne, Surgery impacts right atrial function in tetralogy of Fallot, *J. Thorac. Cardiovasc. Surg.* 147 (4) (2014) 1306–1311.
- [27] S. Cakal, E. Eroglu, O. Baydar, B. Cakal, M.V. Yazicioglu, M. Bulut, C. Dundar, K. Tigen, B. Ozkan, G. Acar, A. Avci, A.M. Esen, Two-dimensional strain and strain rate imaging of the left atrium and left ventricle in adult patients with atrial septal defects before and after the later stage of percutaneous device closure, *Echocardiography* 32 (2014) 470–474 (Mount Kisco, N.Y.).
- [28] B. Ren, L.E. de Groot-de Laat, M.L. Geleijnse, Left atrial function in patients with mitral valve regurgitation, *Am. J. Physiol. Heart Circ. Physiol.* 307 (10) (2014) H1430–7.
- [29] K. Inai, T. Nakanishi, Y. Mori, H. Tomimatsu, M. Nakazawa, Left ventricular diastolic dysfunction in Ebstein's anomaly, *Am. J. Cardiol.* 93 (2) (2004) 255–258.
- [30] S. Kutty, A. Padiyath, L. Li, Q. Peng, S. Rangamani, A. Schuster, D.A. Danford, Functional maturation of left and right atrial systolic and diastolic performance in infants, children, and adolescents, *J. Am. Soc. Echocardiogr.* 26 (4) (2013) 398–409.e2.